

Rotating Frames in *SRT*: Sagnac's Effect and Related Issues*

W. A. Rodrigues, Jr.^{(1,2)†} and M. Sharif^{(1)‡}

⁽¹⁾Institute of Mathematics, Statistics and Scientific Computation

IMECC-UNICAMP CP 6065

13083-970 Campinas, SP, Brazil

and

⁽²⁾Wernher von Braun

Advanced Research Center, UNISAL

Av. A. Garret, 257

13087-290 Campinas, SP Brazil

pacs: 04.90+e 03.30+p

05/10/2001

Contents

1	Introduction	2
2	Some Basic Definitions	3
3	Rotating Reference Frames in <i>SRT</i>	4
3.1	Sagnac Effect	6
3.2	Analysis of Claims by Chiu, Hsu and Sherry	9
3.3	Another <i>nacs</i> to the Frame of a Rotating Platform.	10
4	Conclusions	11
A	Appendix A	11
B	Appendix B	12

*published in *Found. Phys.* **31**, 1767-1783 (2001).

†e-mail: walrod@ime.unicamp.br

‡Permanent Address: Department of Mathematics, Punjab University, Quaid-e-Azam Campus Lahore-54590, PAKISTAN, e-mail: hasharif@yahoo.com

Abstract

After recalling the rigorous mathematical representations in Relativity Theory (*RT*) of (i): observers, (ii): reference frames fields, (iii): their classifications, (iv) naturally adapted coordinate systems (*nacs*) to a given reference frame, (v): synchronization procedure and some other key concepts, we analyze three problems concerning experiments on rotating frames which even now (after almost a century from the birth of *RT*) are sources of misunderstandings and misconceptions. The first problem, which serves to illustrate the power of rigorous mathematical methods in *RT* is the explanation of the Sagnac effect (*SE*). This presentation is opportune because recently there are many non sequitur claims in the literature stating that the *SE* cannot be explained by *SRT*, even disproving this theory or that the explanation of the effect requires a new theory of electrodynamics. The second example has to do with the measurement of the one way velocity of light in rotating reference frames, a problem for which many wrong statements appear in recent literature. The third problem has to do with claims that only Lorentz like type transformations can be used between the *nacs* associated to a reference frame mathematically modeling of a rotating platform and the *nacs* associated with a inertial frame (the laboratory). We show that these claims are equivocated.

1 Introduction

In this paper after recalling the rigorous mathematical definitions of observers, reference frames, naturally adapted coordinate system (*nacs*) to a given reference frame, the synchronization procedure and some other related issues according to the formalism of *RT* we show that: (i) Contrary to recent claims [1-6], Sagnac's effect [7] has a trivial explanation within *RT*¹; (ii) We discuss also some problems connected with the standard synchronization of standard clocks at rest on reference frames in roto-translational motion and what can be expected for the measurement of the one way velocity of light in such a frame. We show that our results are at variance with statements found e.g., in [1,9]; (iii) We investigate also claims [16-19] that the naturally adapted coordinate system (*nacs*) to the reference frame modelling a rotating platform must be related with the *nacs* associated with an inertial frame modelling the laboratory by a Lorentz like type transformation [see eq.(ref31) later], showing that these claims are without any physical or mathematical meaning.

¹This fact is clear from the the very well known paper by Post [8] and also from the more recent references [10-14]. Even if all these references are very good we think that there is space for a new presentation of the subject using the modern concept of *reference frames fields*, because such a presentation permit us to clarify some other obscure issues concerning rotating reference systems (see sections 3.1 and 3.2).

2 Some Basic Definitions

Let τ be a spacetime theory as defined in [19, 24]. Let $Mod\tau$ be the class of all models of τ . Theory τ is said to be relativistic theory if each $\tau \in Mod\tau$ contains a substructure $ST = \langle M, D, \mathbf{g} \rangle$ that is relativistic spacetime as defined in [19,24]. We remember here that $\mathbf{g}$ is a Lorentz metric and D is the Levi-Civita connection of $\mathbf{g}$ on M .

Definition 1. Let ST be a relativistic spacetime. A *moving frame* at $x \in M$ is a basis for the tangent space $T_x M$. An *orthonormal frame* for $x \in M$ is a basis of orthonormal vectors for $T_x M$.

Proposition 1. Let $\mathbf{Q} \in \sec TU \subset \sec TM$ be a time-like vector field such that $\mathbf{g}(\mathbf{Q}, \mathbf{Q}) = 1$. Then, there exist, in a coordinate neighborhood U , three space-like vector fields which together with $\mathbf{Q}$ form an orthogonal moving frame for $x \in U$. The proof is trivial [25].

Definition 2. A non-spinning particle on ST is a pair (m, σ) where $\sigma : \mathcal{R} \supset I \rightarrow M$ is a future pointing causal curve [19-22] and $m \in [0, +\infty)$ is the mass. When $m = 0$ the particle is called a photon. When $m \in (0, +\infty)$ the particle is said to be a material particle. σ is said to be the world line of the particle.

Definition 3. An *observer* in $\langle M, D, \mathbf{g} \rangle$ is a future pointing time-like curve $\gamma : \mathcal{R} \supset I \rightarrow M$ such that $\mathbf{g}(\gamma_* u, \gamma_* u) = 1$. The inclusion parameter $I \rightarrow \mathcal{R}$ in this case is called the proper time along γ , which is said to be the world line of the observer.

Observation 1. The physical meaning of proper time is discussed in details, e.g., in [20,21] which deals with the theory of time in relativistic theories.

Definition 4. An *instantaneous observer* is an element of TM , i.e., a pair $(z, \mathcal{Z})$, where $z \in M$, and $\mathcal{Z} \in T_z M$ is a future pointing unit time-like vector.

The Proposition 1 together with the above definitions suggests:

Definition 5. A *reference frame* in $ST = \langle M, D, \mathbf{g} \rangle$ is a time-like vector field which is a section of $TU, U \subseteq M$ such that each one of its integral lines is an observer.

Observation 2. In [19,22] an arbitrary reference frame $\mathbf{Q} \in \sec TU \subseteq \sec TM$ is classified as (i), (ii) below.

(i) according to its *synchronizability*. Let $\alpha_{\mathbf{Q}} = g(\mathbf{Q}, \cdot)$. We say that $\mathbf{Q}$ is locally synchronizable iff $\alpha_{\mathbf{Q}} \wedge d\alpha_{\mathbf{Q}} = 0$. $\mathbf{Q}$ is said to be locally proper time synchronizable iff $d\alpha_{\mathbf{Q}} = 0$. $\mathbf{Q}$ is said to be synchronizable iff there are C^∞ functions $h, t : M \rightarrow \mathcal{R}$ such that $\alpha_{\mathbf{Q}} = hdt$ and $h > 0$. These definitions are intuitive.

(ii) according to the *decomposition*² of

$$D\alpha_{\mathbf{Q}} = \mathbf{a}_{\mathbf{Q}} \otimes \alpha_{\mathbf{Q}} + \omega_{\mathbf{Q}} + \sigma_{\mathbf{Q}} + \frac{1}{3}\Theta_{\mathbf{Q}}\mathbf{h}, \quad (1)$$

where

$$\mathbf{h} = \mathbf{g} - \alpha_{\mathbf{Q}} \otimes \alpha_{\mathbf{Q}} \quad (2)$$

²The validity of decomposition (1) is proved in Appendix C.

is called the projection tensor (and gives the metric of the rest space of an instantaneous observer [20,22]), $\mathbf{a}_Q$ is the (form) acceleration of Q , ω_Q is the rotation of Q , σ_Q is the shear of Q and Θ_Q is the expansion of Q . In a coordinate chart (U, x^μ) , writing $Q = Q^\mu \partial / \partial x^\mu$ and $\mathbf{h} = (g_{\mu\nu} - Q_\mu Q_\nu) dx^\mu \otimes dx^\nu$ we have

$$\begin{aligned}\omega_{Q\mu\nu} &= Q_{[\mu;\nu]}, \\ \sigma_{Q\alpha\beta} &= [Q_{(\mu;\nu)} - \frac{1}{3}\Theta_Q h_{\mu\nu}] h_\alpha^\mu h_\beta^\nu, \\ \Theta_Q &= Q^\mu{}_{;\mu}.\end{aligned}\tag{3}$$

We shall need in what follows the following result that can be easily proved:

$$\alpha_Q \wedge d\alpha_Q = 0 \Leftrightarrow \omega_Q = 0.\tag{4}$$

Eq.(3) means that rotating reference frames (i.e., frames for which $\omega_Q \neq 0$) are not locally synchronizable and this result is the key in order to solve the misconceptions usually associated with rotating reference frames as the ones studied in section 3.

Observation 3. In Special Relativity where the space time manifold is $< M = \mathcal{R}^4, \mathbf{g} = \eta, D^\eta >$ ³ an *inertial reference frame* (IRF) $\mathbf{I} \in \text{sec}TM$ is defined by $D^\eta \mathbf{I} = 0$.

Observation 4. Before concluding this section it is very much important to recall that a reference frame field as introduced above is a *mathematical* instrument. It did not necessarily need to have a material substratum (i.e., to be realized as a material physical system) in the points of the spacetime manifold where it is defined⁴. More properly, we state that the integral lines of the vector field representing a given reference frame do not need to correspond to worldlines of real particles. If this crucial aspect is not taken into account we may incur in serious misunderstandings. This observation will become clear in what follows.

3 Rotating Reference Frames in *SRT*

In order to grasp the use of the very abstract notions introduced above we shall study in this sections three problems. In section 3.1 we give an analysis of the Sagnac effect [7], since there are many claims in the recent literature saying that this effect cannot be explained by *SRT* [1-3] or even requires the formulation of a new electrodynamics where the photon has a non zero mass [4]⁵ or where the usual $U(1)$ gauge theory is substituted by a $SU(2)$ [5] (or $O(3)$, [6]) gauge

³ η is a constant metric, i.e., there exists a chart $\langle x^\mu \rangle$ of $M = \mathcal{R}^4$ such that $\eta(\partial/\partial x^\mu, \partial/\partial x^\nu) = \eta_{\mu\nu}$, the numbers $\eta_{\mu\nu}$ forming a diagonal matrix with entries $(1, -1, -1, -1)$. Also, D^η is the Levi-Civita connection of η .

⁴More on this issue is discuted in [24].

⁵Vigier [4] also claims that the Sagnac effect implies in the existence of a fundamental frame.

theory⁶. In section 3.2 we analyse the claims of [9] that supposedly proved that clocks on a rotating platform that are synchronized à l'Einstein at a given instant will not measure the standard value of the *one way* velocity of light after that the platform makes half a rotation (as seen by an *IRF*). In section 3.3 we analyze the claims of [17-18] that it is necessary in order to describe a rotating platform, coordinate functions necessarily different from the coordinate functions defined by eq.(11) later.

In this section $(M, \mathbf{g}, D)$ denotes Minkowski spacetime and let $\mathbf{I} \in \sec TM$ be an inertial reference frame on M . Let (t, r, ϕ, z) be cylindrical coordinates for M such that $\mathbf{I} = \partial/\partial t$. Then

$$\mathbf{g} = dt \otimes dt - dr \otimes dr - r^2 d\phi \otimes d\phi - dz \otimes dz \quad (5)$$

Let $\mathbf{P} \in TM$ be another reference frame on M given by

$$\mathbf{P} = (1 - \omega^2 r^2)^{-1/2} \frac{\partial}{\partial t} + \omega r^2 (1 - \omega^2 r^2)^{-1/2} \frac{\partial}{\partial \phi} \quad (6)$$

$\mathbf{P}$ is well defined in an open set $U \subset M$ such that (Note that the coordinate system (t, r, ϕ, z) is not submitted to the restriction $0 < r < 1/\omega$ of Eq.(9))

$$U \equiv (-\infty < t < \infty; 0 < r < 1/\omega; 0 \leq \phi < 2\pi; -\infty < z < \infty) \quad (7)$$

Then,

$$\alpha_{\mathbf{P}} = \mathbf{g}(\mathbf{P}, \cdot) = (1 - \omega^2 r^2)^{-1/2} dt - \omega (1 - \omega^2 r^2)^{-1/2} d\phi \quad (8)$$

and

$$\alpha_{\mathbf{P}} \wedge d\alpha_{\mathbf{P}} = -\frac{2\omega r}{(1 - \omega^2 r^2)^2} dt \wedge dr \wedge d\phi \quad (9)$$

The rotation (or vortex) vector $[\Omega = \tilde{\mathbf{g}}(*(\alpha_{\mathbf{P}} \wedge d\alpha_{\mathbf{P}}),)]$ is

$$\Omega = \omega (1 - \omega^2 r^2)^{-1/2} \frac{\partial}{\partial z}, \quad (10)$$

where $\tilde{\mathbf{g}}$ is the metric in the cotangent bundle. This means that $\mathbf{P}$ is rotating with angular velocity ω (as measured) according to $\mathbf{I}$ in the z direction and local angular velocity $\omega/(1 - \omega^2 r^2)^{-1/2}$ as measured by an observer at rest on $\mathbf{P}$ at a point with coordinates $(t, r, \phi, z = 0)$. This result is clear if we remember that a clock at rest on $\mathbf{P}$ at $(t, r, \phi, z = 0)$ has its period greater by a factor $(1 - \omega^2 r^2)^{-1/2}$ in relation to clocks at rest on $\mathbf{I}$.

Now $\mathbf{P}$ can be realized on $U \subset M$ by a rotating platform, but is obvious that at the same 'time' on U , $\mathbf{I}$ cannot be realized by any physical system.

$\mathbf{P}$ is a typical example of a reference frame for which it does not exist a $(nacs|\mathbf{P})$ such that the time like coordinate of the frame has the meaning of proper time registered by standard clocks at rest on $\mathbf{P}$.

⁶The wrong claims of [1-3] have also been discussed and clarified in two recent papers by Tartaglia and Rizzi [13] and by Matolcsi [14]. We decided to present our analysis of the claims of [1-3] because we think that our presentation has complementary material concerning the ones in [11,12] and because it serves the proposal of showing how the concept of reference frames as introduced here works. Moreover, we are not going to discuss the claims of [4] and of [5,6] in this paper, leaving such an analysis for a new paper. Here, we only call the reader's attention that the proposal of [6] is almost identical to that of [5], which could eventually be a difficult reference to be found.

3.1 Sagnac Effect

According to the classification of reference frames [20-22] it is indeed trivial to see that $\mathbf{P}$ is *not* proper time synchronizable or even locally synchronizable.

This fact is *not* very well known by the majority of physicists (which are not specialists in RT) and leads people from time to time to claim that optical experiments done on a rotating platform *disproves SRT*. A recent claim of this kind has been done by Selleri and collaborators on a series of papers [1-3] and Vigier [4] who thinks that in order to explain the effects observed it is necessary to attribute a non-zero mass to the photon and that this implies in the existence of a fundamental reference frame.

There are also some papers that claim that in order to describe the Sagnac effect it is necessary to build a non-abelian electrodynamics [5,6]. This is only a small sample of papers containing wrong statements concerning the Sagnac experiment. Now, the Sagnac effect is a well established fact (used in the technology of the gyro-ring) that the transit time employed for a light ray to go around a closed path enclosing a non-null area depends on the sense of the curve followed by the light ray.

Selleri arguments that such a fact implies that the velocity of light as determined by observers on the rotating platform cannot be constant, and must depend on the direction, implying that nature realizes a synchronization of the clocks which are at rest on the platform different from the one given by the Einstein method.

Selleri arguments also that each small segment of the periphery of the rotating platform of radius R can be thought as at rest on an inertial reference frame moving with speed ωR relative to the laboratory (here modeled by $\mathbf{I}$).

In that way, if the synchronization is done à l'Einstein between two clocks at neighboring points of the small segment, the resulting measured value of the one way velocity of light must result constant in both directions ($c = 1$), thus contradicting the empirical fact demonstrated by the Sagnac effect.

Now, we have already said that $\mathbf{P}$ is not proper time synchronizable, nor is $\mathbf{P}$ locally synchronizable (in general), as follows from the rigorous theory of time in Relativity [20-22].

However, for two neighboring clocks at rest on the periphery of a uniformly rotating platform an Einstein's synchronization can be done. Let us see what we get.

First, let $\langle \hat{x}^\mu \rangle$ be a $(nacs|\mathbf{P})$ given by

$$\hat{t} = t, \quad \hat{r} = r, \quad \hat{\phi} = \phi - \omega t, \quad \hat{z} = z \quad (11)$$

In these coordinates $\mathbf{g}$ is written as

$$\begin{aligned} \mathbf{g} &= (1 - \omega^2 \hat{r}^2) d\hat{t} \otimes d\hat{t} - 2\omega \hat{r}^2 d\hat{\phi} \otimes d\hat{t} - d\hat{r} \otimes d\hat{r} - r^2 d\hat{\phi} \otimes d\hat{\phi} - d\hat{z} \otimes d\hat{z} \\ &= g_{\mu\nu} dx^\mu \otimes dx^\nu \end{aligned} \quad (12)$$

Now take the two clocks A and B, at rest on $\mathbf{P}$. Suppose they follow the world lines ρ and ρ' which are infinitely close.

According to the chronometric postulate (see, e.g., [21,22]) the events on ρ or ρ' can be orderly. We write, e.g., as usual $e_1 < e < e_2$ to indicate that on ρ , e is posterior to e_1 and e_2 is posterior to e .

Now let

$$e_1 = (x_{e_1}^0, x_{e_1}^1, x_{e_1}^2, x_{e_1}^3), \quad e_2 = (x_{e_2}^0, x_{e_2}^1, x_{e_2}^2, x_{e_2}^3),$$

$$e' = (x_{e'}^0, x_{e'}^1, x_{e'}^2, x_{e'}^3), \quad e = (x_e^0, x_e^1 + \Delta x^1, x_e^2 + \Delta x^2, x_e^3 + \Delta x^3), \quad (13)$$

e_1 is the event on ρ when a light signal is sent from clock A to clock B. e' is the event when the light signal arrives at clock B on ρ' and is (instantaneously) reflected back to clock A where it arrives at event e_2 . e is the event simultaneously on ρ to the event e' according to Einstein's convention and we have [20,21,26]

$$x_e^0 = x_{e'}^0 + \frac{g_{0i}}{g_{00}} \Delta x^i \neq x_{e'}^0 \quad (14)$$

We emphasize that Eq.(14) does not mean that we achieved a process permitting the synchronization of two standard clocks following the world lines ρ and ρ' , *because* standard clocks in general do not register the “flow” of the time-like coordinate x^0 . However, in some particular cases such as when $g_{\mu\nu}$ is independent of time and for the specific case where the clocks are very near (see below) and at rest on the periphery of a uniformly rotating platform this can be done.

This is so because standard clocks at rest at the periphery of a uniformly rotating platform “tick tack” at the same ratio relative to **I**. Once synchronized they will *remain* synchronized. It follows that the velocity of light measured by these two clocks will be independent of the direction followed by the light signal and will result to be $c = 1$ every time that the measurement will be done. This statement can be *trivially* verified and is in complete disagreement with a proposal of [9] that we discuss in section 3.2. We now analyze with more details what will happen if we try the *impossible* task (since **P** is not proper time synchronizable, as already said above) of synchronizing standard clocks at rest on a rotating platform which is the material support of the reference frame **P**.

Suppose that we synchronize (two by two) a series of standard clocks (such that any two are very close⁷) at rest and living on a closed curve along the periphery of a rotating platform. Let us number the clocks as 0, 1, 2, ..., n . Clocks 0 and 1 are supposed “be” at the same point p_1 and are the beginning of our experiment synchronized. After that we synchronize, clock 1 with 2, 2 with 3, ... and finally n with 0. From Eq.(14) we get immediately that at the end of the experiment clocks 0 and 1 will not be synchronized and the coordinate time difference between them will be

$$\Delta \hat{t} = - \oint \frac{g_{0i}}{g_{00}} d\hat{x}^i = \oint \frac{\omega R^2}{1 - \omega^2 R^2} d\hat{\phi} \quad (15)$$

⁷Very close means that $l/R \ll 1$, where l is the distance between the clocks and R is the radius of the platform, both distances being done in the frame **P**.

For $\omega R \ll 1$ we have $\Delta\hat{t} = \pm 2\omega S$ where S is the area of the rotating platform and the signals $\pm$ refer to the two possible directions in each we can follow around the rotating platform.

The correct relativistic explanation of the Sagnac experiment is as follows. Suppose (accepting the validity of the geometrical optics approximation [8]) that the world line of a light signal that follows the periphery of the rotating platform of radius R is the curve $\sigma : \mathcal{R} \supset I \rightarrow \mathcal{M}$ such that σ_* is a null vector. Using the coordinate $\hat{t}$ as a curve parameter we have

$$\mathbf{g}(\sigma_*, \sigma_*) = (1 - \omega^2 R^2)(d\hat{t} \circ \sigma)^2 - 2\omega R^2(d\hat{\phi} \circ \sigma)(d\hat{t} \circ \sigma) - R^2(d\hat{\phi} \circ \sigma)^2 = 0 \quad (16)$$

Then

$$\frac{d\hat{\phi} \circ \sigma}{d\hat{t}} = -\omega \pm 1/R \quad (17)$$

Then, the coordinate times for a complete round are

$$\hat{T}_{\pm} = 2\pi R / 1 \mp \omega R \quad (18)$$

where the signals $\pm$ refer to the two possible paths around the periphery, with $-$ when the signal goes in the direction of rotation and $+$ in the other case. It is quite obvious that $\hat{T}_{\pm}$ can be measured by a single clock.

Now, observe that the length⁸ of the periphery as measured with a set of standard rulers at rest on a rotating platform is

$$L = 2\pi R(1 - \omega^2 R^2)^{-1/2}. \quad (19)$$

Being $\tau_{\pm}$ the proper times measured by standard clock at rest at the periphery of the rotating platform, corresponding to $\hat{T}_{\pm}$, we have

$$\tau_{\pm} = (1 - \omega^2 R^2)^{1/2} \hat{T}_{\pm} = L(1 \pm \omega R). \quad (20)$$

This equation explains trivially the Sagnac effect according to Special Relativity.

Selleri [1] calls the quantities $c_{\pm}$

$$\frac{1}{c_{\pm}} = \frac{L}{\tau_{\pm}} \quad (21)$$

the global velocities of light in the rotating platform for motions of light in the directions of rotation ($-$) and in the contrary sense ($+$). He then argues that these values must also be the local values of the *one-way* velocities of light, i.e., the values that an observer would necessarily measure for light going from a point p_1 in the periphery of a rotating platform to a neighboring point p_2 . He even believes to have presented an ontological argument that implies that

⁸This result follows at once from the definition of the projection tensor $\mathbf{h}$ (eq.(2)) which gives the metric of an instantaneous observer. The length L of the periphery given by eq.(19) is just the sum of the measurements done by different instantaneous observers along the periphery of the disc. Note that the claim by Klauber [15] that the space geometry of the disc is flat is in contradiction with the way measurements are realized in RT .

Special Relativity is not true. Well, what is wrong with Selleri's argument is the following. Although it is true that the global velocities $c_{\pm}$ can be measured with a single clock, the measurement of the local one way velocity of light transiting between p_1 and p_2 requires two standard clocks synchronized at that points. Local Einstein synchronization is possible and as described above gives a local velocity equal to $c = 1$, and this fact leads to no contradiction.

We observe at this point that the claims of [1-3] have also been discussed properly by [13-14], and our analysis touch on some complementary aspects of the problem. The analysis of [14] is specially rigorous and elegant being based on an intrinsic approach to RT which does not use reference frames and that has been developed in detail in [27].

3.2 Analysis of Claims by Chiu, Hsu and Sherry

Another *interesting* result, worth to be mentioned here because it is a source of a misunderstanding is the following one that appears in a paper by Chiu, Hsu and Sherry [9].

Take two clocks A and B (very close, i.e., as above $l/R \ll 1$) and at rest at the periphery of a rotating platform ($\mathbf{P}$ frame) with center at the point with space coordinates $(0, R, 0)$ according to an $\mathbf{I}$ frame (the laboratory). Let

$$\mathbf{I}' = \frac{1}{\sqrt{1 - \omega^2 R^2}} \partial/\partial t + \frac{\omega R}{\sqrt{1 - \omega^2 R^2}} \partial/\partial x \quad (22)$$

be an inertial Lorentz frame moving with speed ωR relative to $\mathbf{I}$ and such that at the beginning of the synchronization procedure, clock A is at rest relative to $\mathbf{I}'$ and clock B is differentially at rest relative to $\mathbf{I}'$. It is assumed that the platform is uniformly rotating in the xy-plane and clocks A and B are synchronized when A is at the point with coordinates $(0, -\varepsilon, 0, 0)$ according to $\mathbf{I}$ and $(0, -(1 - \omega^2 R^2)^{-\frac{1}{2}} \varepsilon, 0, 0)$ according to $\mathbf{I}'$. Of course, clocks synchronized in that way will not be synchronized according to the point of view of an observer in the $\mathbf{I}''$ frame,

$$\mathbf{I}'' = \frac{1}{\sqrt{1 - \omega^2 R^2}} \partial/\partial t - \frac{\omega R}{\sqrt{1 - \omega^2 R^2}} \partial/\partial x. \quad (23)$$

This means that when the two clocks A and B are such that the coordinates of A is $(0, \varepsilon, 2R, 0)$ according to $\mathbf{I}$, an observer at rest in the $\mathbf{I}''$ frame comoving with the clock A and such that at that event clock B is also at rest will see the clocks out of phase if compared with their clocks that have been synchronized à l'Einstein. This is so because $\mathbf{I}''$ is moving relative to $\mathbf{I}$, and so clocks which are synchronized relative to $\mathbf{I}'$ will not be synchronized according to $\mathbf{I}''$. Now authors of [9] suppose that each segment (around the equator of Earth) can be thought as at rest in an instantaneous inertial frame. Based on this argument they claim that the velocity of light measured by the stable clocks A and B synchronized as above described will suffer diurnal variations. Of course, this is *not* true. The observers at rest on Earth will continue to find $c = 1$ as the

result of the measured one-way velocity of light. And before leaving this section it is important to emphasize that the supposition that a segment of the equator of Earth (or the periphery of a rotating platform) is instantaneously at rest on an inertial comoving frame, say $\mathbf{I}'$, do not transform this segment in a part of that frame. It continues to be part of the rotating frame, say $\mathbf{P}$ and observers living in $\mathbf{P}$ can always know that they are rotating, since rotation is an absolute concept in RT .

We take the opportunity to call the reader's attention that the non synchronizability of clocks on a rotating reference frame is crucial for an understanding of the time as recorded by satellites (the GPS) and the time registered by clocks at rest on the Earth, as described by Scorgie [28]. On this issue see also [29].

Also, it is important to kept in mind that although we have speaked and worked only with Einstein's synchronization in the above analysis, non standard synchronizations can be also used. References on this important issue are [30-35]

3.3 Another *nacs* to the Frame of a Rotating Platform.

Some authors, e.g., [16-18] start the discussion of the rotating platform problem by supposing that the *natural* coordinate functions adapted to a rotating platform must *necessarily* be related with a (*nacs*| $\mathbf{I}$) by the following Lorentz like type transformations:

$$\begin{aligned}\bar{t} &= t \cosh \Omega r - r \phi \sinh \Omega r, \\ \bar{r} &= r, \quad \bar{z} = z. \\ \bar{\phi} &= \phi \cosh \Omega r - \frac{t}{r} \sinh \Omega r,\end{aligned}\tag{24}$$

These authors claim that these transformations are necessary and the authors of [19] even claim that such coordinates are necessary to solve some puzzles concerning the partial polarization of electrons (or positrons) in a storage ring, for which they think to have found an explanation by appealing to the Unruh-Davies effects that according to quantum field theory must exist for some kinds of accelerated motion. We defer the discussion of the last problem to another paper. Here we are interested in a much simple problem: which is the reference frame field $\bar{\mathbf{P}}$ such that the coordinates $(\bar{t}, \bar{r}, \bar{\phi}, \bar{z})$ are a (*nacs*— $\mathbf{P}$)? The answer is:

$$\bar{\mathbf{P}} = (\cosh \Omega r) \partial / \partial t - \left(\frac{1}{r} \sinh \Omega r \right) \partial / \partial \phi = \partial / \partial \bar{t}.\tag{25}$$

Now, if we impose that $\bar{\mathbf{P}}$ is realized by a physical system (as it is the case of the $\mathbf{P}$ frame) then the velocity of the periphery of the platform with radius $r = \bar{R}$, must satisfy the following constrain

$$\lim_{r \rightarrow \bar{R}} \sinh \Omega r = 1.\tag{26}$$

Now it is trivial to verify that $\alpha_{\bar{\mathbf{P}}} \wedge d\alpha_{\bar{\mathbf{P}}} \neq 0$ and then $\bar{\mathbf{P}}$ (like $\mathbf{P}$) is not proper time synchronizable. Moreover, the other characteristics of the $\bar{\mathbf{P}}$ frame [given

by the *unique* decomposition showed in eq.(1)] will be the same as for the $\bar{\mathbf{P}}$ frame if

$$\Omega = \frac{1}{r} \operatorname{arcsinh} \omega \frac{r}{\sqrt{(1 - \omega^2 r^2)}} \quad (27)$$

In fact, with this identification $\bar{\mathbf{P}} \equiv \mathbf{P}$ and this, of course is the only satisfactory solution for our problem. Moreover, this shows that to think that a Lorentz like type transformation [eq.(24)] has some special significance is a nonsequitur and to think that these coordinates help to solve the polarization problem of electrons in storage rings is to forget that coordinates are labels, not the physics and to mistaken the concept of coordinates charts with the concept of frames. In *RT* we can use any *arbitrary* coordinate chart to do calculations, but in that theory as well as in quantum field theory, in general different reference frames are not *physically equivalent*, a concept that we study in [24].

4 Conclusions

In this paper we review some crucial issues associated with the concept of a reference frame in *SRT*. By properly defining this concept and others associated with it we investigated some issues concerning rotating reference frames in Minkowski spacetime. We apply consistently the mathematical notions of section 1, first to the study of the Sagnac effect in *SRT*. This has been done because it has been argued recently once again by different people that Sagnac's effect cannot be explained by *SRT*. We disprove this statement by showing that this effect has a simple explanation in that theory. Further we explore the problems which arise in synchronization of standard clocks at rest on a rotating platform and the problem of the measurement of the one way velocity of light in that frames, clarifying many misconceptions that also appeared in recent literature. We discuss also some claims that a more appropriate coordinate system [than the canonical one given by eq.(11)] adapted to the frame that mathematically models an uniforming rotating platform is given by a Lorentz like transformation [eq.(24)] between the coordinates of the frame and the *(nacs|I)*, where **I** is the inertial frame modelling the laboratory. We show that such claims are equivocated.

A Appendix A

In this appendix we show the reason for the definition of a synchronizable frame.

(i) We recall that a frame $\mathbf{Q} \in \sec TM$ is said to be synchronizable iff there are C^∞ functions $h, \bar{t}: M \rightarrow \mathfrak{R}$ such that $\alpha_Q = h d\bar{t}$ and $h > 0$.

(ii) Now, we need to construct a proof for Proposition 1. Suppose that the metric of the manifold in the chart (U, η) with coordinate functions $\langle x^\mu \rangle$ is $g = g_{\mu\nu} dx^\mu \otimes dx^\nu$. Let $Z = (Z^\mu \partial / \partial x^\mu) \in \sec TM$ be an arbitrary reference frame and $\alpha_Z = g(Z, \cdot) = Z_\mu dx^\mu$, $Z_\mu = g_{\mu\nu} Z^\nu$. Then, $g_{\mu\nu}(x) Z^\mu Z^\nu = 1$.

Now, define

$$\begin{aligned}\theta^0 &= (\alpha_Z)_\mu dx^\mu = Z_\mu dx^\mu, \\ \gamma_{\mu\nu} &= g_{\mu\nu} - Z_\mu Z_\nu;\end{aligned}\tag{A1}$$

Then the metric g can be written due to the hyperbolicity of the manifold as

$$\begin{aligned}g &= \theta^0 \otimes \theta^0 - \sum_{i=1}^3 \theta^i \otimes \theta^i \\ \sum_{i=1}^3 \theta^i \otimes \theta^i &= \gamma_{\mu\nu}(x) dx^\mu \otimes dx^\nu\end{aligned}\tag{A2}$$

(iii) Now, suppose that there exist a coordinate chart (V, χ) , $U \cap V \neq \emptyset$ with coordinate functions $\langle \bar{x}^\mu \rangle$ such that $Z = \frac{1}{\sqrt{g_{00}}} \partial / \partial \bar{x}^0$. Then we can verify that

$$\theta^0 = \sqrt{g_{00}(\bar{x})} d\bar{x}^0 + \frac{g_{i0}(\bar{x})}{g_{00}(\bar{x})} d\bar{x}^i,\tag{A3}$$

$$\begin{aligned}\sum_{i=1}^3 \theta^i \otimes \theta^i &= \gamma_{\mu\nu}(\bar{x}) d\bar{x}^\mu \otimes d\bar{x}^\nu \\ &= \left(\frac{g_{i0}(\bar{x})g_{j0}(\bar{x})}{g_{00}(\bar{x})} - g_{ij}(\bar{x}) \right) d\bar{x}^i \otimes d\bar{x}^j, \quad i, j = 1, 2, 3.\end{aligned}\tag{28}$$

Now let us apply the results of (ii) and (iii) above a synchronizable frame $\mathbf{Q}$ defined in (i). In the coordinate chart (V, χ) with the coordinate functions $\langle \bar{x}^\mu \rangle$ we take $\bar{x}^0 = \bar{t}$. Since in these coordinates $\alpha_Q = h d\bar{t}$, it follows that in these coordinates $\mathbf{Q} = \frac{1}{h} \partial / \partial \bar{x}^0$ and $g_{00}(\bar{x}) = h^2$.

From the first line of Eq.(A3) it follows that in this case $g_{i0}(\bar{x}) = 0$, i.e., the metric in the coordinates $\langle \bar{x}^\mu \rangle$ is diagonal.

(iv) Now, recall Eq.(14) from the text that gives the time like coordinates of two events e and e' that are simultaneous according to Einstein's simultaneity definition. We see that in the coordinates $\langle \bar{x}^\mu \rangle$ the events e and e' are such that $\bar{x}_e^0 = \bar{x}_{e'}^0$. This condition justify the definition of a synchronizable reference frame and turns the definition a very intuitive indeed.

B Appendix B

Here we explain the meaning of the definition of a locally synchronizable reference frame $\mathbf{Q} \in \sec TM$, for which $\alpha_Q \wedge d\alpha_Q = 0$.

We recall the following:

(i) A 3-direction vector field H_x in a 4-dimensional manifold M for $x \in M$ is a 3-dimensional vector subspace H_x of $T_x M$ which satisfies a *differentiability condition*. This condition is usually expressed as:

(dfc1) For each point $x_0 \in M$ there is a neighborhood $U \subset M$ of x_0 such that there exist three differential vector fields, $X_i \in \sec TM$ ($i = 1, 2, 3$) such that $X_1(x), X_2(x), X_3(x)$, is a basis of $H_x, \forall x \in U$, or:

(dfc2) For each point $x_0 \in M$ there is a neighborhood $U \subset M$ of x_0 such that there exist a one form $\alpha \in \sec T^*M$ such that for all $X \in H_x \Leftrightarrow \alpha(X) = 0$.

(ii) We now recall Frobenius theorem [36]: Let be $x_0 \in M$ and let be U a neighborhood of x_0 . In order that, for each $x_0 \in M$ there exist a 3-dimensional manifold $\Pi_{x_0} \ni x_0$ (called the integral manifold through x_0) of the neighborhood U , *tangent* to H_x for all $x \in \Pi_{x_0}$ it is necessary and sufficient that $[X_i, X_j]_x \in H_x$, (for all $i, j = 1, 2, 3$ and $\forall x \in \Pi_{x_0}$) if we consider condition (dfc1). If we consider the (dfc2) then a necessary and sufficient condition for the existence of Π_{x_0} is that

$$\alpha \wedge d\alpha = 0. \quad (\text{B1})$$

(iii) Now, let us apply Frobenius theorem for the case of a Lorentzian manifold. Let $\mathbf{Q} \in \sec TM$ be a reference frame for which $\alpha_{\mathbf{Q}} \wedge d\alpha_{\mathbf{Q}} = 0$.

Then, from the condition for the existence of a integral manifold through $x_0 \in M$ we can write,

$$\alpha_{\mathbf{Q}}(X_1) = \alpha_{\mathbf{Q}}(X_2) = \alpha_{\mathbf{Q}}(X_3) = 0. \quad (\text{B2})$$

Now, since $\mathbf{Q}$ is a time like vector field, Eq.(B2) implies that the X_i ($i = 1, 2, 3$) are spacelike vector fields. It follows that the vector field $\mathbf{Q}$ is orthogonal to the integral manifold Π_{x_0} which is in this case a spacelike surface.

Now the meaning of a synchronizable reference frame (which is given by Eq.(B1) becomes clear. Observers in such a frame can locally separate any neighborhood U of x_0 (where they “are”) in time $\times$ space.

C Appendix C

In this paper we prove the decomposition of a general reference frame $\mathbf{Q} = (Q^\mu \partial / \partial x^\mu) \in \sec TM$ given by Eq.(1) in the text.

To prove the decomposition, choose a chart (U, η) with coordinate functions $\langle x^\mu \rangle$ and write

$$D\alpha_{\mathbf{Q}} = Q_{\mu;v} dx^\mu \otimes dx^v. \quad (\text{C1})$$

Then, all we need is to verify that the functions $\omega_{\mathbf{Q}_{\mu v}}, \sigma_{\mathbf{Q}_{\mu v}}$ and $\Theta_{\mathbf{Q}}$ given by Eq.(3) of the text satisfy the required conditions and that

$$Q_{\rho;\tau} h_\mu^\rho h_\nu^\tau = Q_{\mu;v} - a_{Q_\mu} Q_\nu, \quad (\text{C2})$$

where in Eq.(C2) h_μ^ρ are the mixt components of the projection tensor $\mathbf{h}$ given by Eq.(2) of the text.

Acknowledgments: One of the authors (MS) would like to thank TWAS for the traveling grant and CNPq for the local hospitality at IMECC-UNICAMP, SP Brazil. Authors are grateful to Professors U. Bartocci, Y. Bozhkov, J. Vaz, Jr. and A. Saa and Drs. A. M. Moya and D. S. Thober for stimulating discussions.

References

- [1] F. Selleri, *Found. Phys.* **26**, 641 (1996); *FPL* **10**, 73 (1997).
- [2] J. Croca J. and F. Selleri, *N. Cimento B* **114**, 447 (1999).
- [3] F. Goy and F. Selleri, *Found. Phys. Lett.* **10**, 17 (1997).

- [4] J. P. Vigiér, *Phys. Lett. A* **234**, 75 (1997).
- [5] T. W. Barrett, *Ann. Fond. L. de Broglie* **14**, 37 (1989); **15**, 143 (1989); **15**, 253 (1990); in A. Lakhatia (ed.), *Essays of the Formal Aspects of Electromagnetic Theory*, World Scientific, Singapore Pub. Co., pp. 6- 86, 1993; in T.W. Barrett and D. M. Grimes (eds.), *Advanced Electromagnetism*, World Scientific, Singapore Pub. Co., pp. 278-313, 1995.
- [6] P. K. Anastasovski, et al (AIAS group), *Found. Phys. Lett.* **12**, 579 (1999).
- [7] M. G. Sagnac, *Comp. Rend.* **507**, 708 (1913); **157**, 1410 (1913).
- [8] E. J. Post, *Rev. Mod. Phys.* **39**, 475 (1967).
- [9] C. B. Chiu, J. P. Hsu and T. N. Sherry, *Phys. Rev. D* **16**, 2420 (1977).
- [10] A. Asthekart and A. Magnon, *J. Math. Phys.* **16**, 341 (1975).
- [11] J. Anadan, *Phys. Rev. D* **24**, 338 (1981).]
- [12] A. A. Logunov, and Yu. V. Chugreev, *Sov. Phys. Usp.* **31**, 861 (1988).
- [13] G. Rizzi and A. Tartaglia, *Found. Phys.* **28**, 1663 (1998); **12**, 179 (1999); **28**, 1663 (1998).
- [14] T. Matolcsi, *Found. Phys.* **28**, 1685 (1998).
- [15] R. D. Klauber, *Found. Phys.* **11**, 405-483 (1998).
- [16] G. Trocheres, *Philos. Mag.* **40**, 1143 (1949).
- [17] Takeno H., *Prog. Theor. Phys.* **7**, 367 (1952).
- [18] V. A. De Lorenzi and N. F. Svaiter, *Found. Physics* **29**, 1233 (1999); *Int. J. Mod. Phys. A* **14**, 717 (1999); in Yu. N. Gnedin, A. A. Grib, V. M. Mostepanenko and W. A. Rodrigues, Jr., (eds.), *Proc. of the Fourth A. Friedmann Int. Symposium on Gravitation and Cosmology*, pp.295-309, IMECC-UNICAMP Publ., Campinas, 1999.
- [19] W. A. Rodrigues, Jr., M. Scanavini and L. P. de Alcantara, *Found. Phys. Lett.* **3**, 59 (1990).
- [20] W. A. Rodrigues, Jr. and M. A. F. Rosa, *Found. Phys.* **19**, 705 (1989).
- [21] W. A. Rodrigues, Jr. and de E.C. Oliveira, *Phys. Lett. A* **140**, 479 (1989).
- [22] R. K. Sachs and H. Wu, *General Relativity for Mathematicians*, Springer Verlag, N. York, Berlin, 1977.
- [23] J. E. Maiorino and W. A. Rodrigues, Jr., What is Superluminal Wave Motion?, *Sci. and Tech. Mag.* **2**(2) (1999), see <http://www.ceptec.br/~stm>.
- [24] W. A. Rodrigues, Jr. and M. Sharif, *Equivalence Principle and the Principle of Local Lorentz Invariance*, *Found. Phys.* **31**(12) (2001).
- [25] Y. Choquet-Bruhat, C. Dewitt-Morette and M. Dillard-Bleick, *Analysis, Manifolds and Physics* (revised version), North Holland Pub. Co., Amsterdam, 1982.
- [26] L. D. Landau and E. M. Lifschitz, *The Classical Theory of Fields* (4th revised english edition), Pergamon Press, Oxford, 1975.
- [27] T. Matolsci, *Spacetime Withouth Reference Frames*, Akadémiai Kiadó, Budapest, 1993.
- [28] G. C. Scorgie, *Eur. J. Phys.* **12**, 64 (1991).
- [29] T. B. Bahder, *IEEE Trans. Aerospace and Electr. Syst.* **34**, 1350 (1998).

- [30] H. Reinchbach, *The Philosophy of Space and Time*, Dover, New York, 1958.
- [31] W. A. Rodrigues, Jr. and J. Tiomno, *Found. Phys.* **15**, 945 (1985).
- [32] R. Mansouri, and R. U. Sexl, *Gen. Rel. Grav.* **8**, 497 (1977).
- [33] R. Anderson, I. Vetharaniam, G. E. Stedman, *Phys. Rep.* **295**, 93 (1998).
- [34] C. Leuber, K. Aufinger, and P. Krumm, *Eur. J. Phys.* **13**, 170 (1992)
- [35] T. Ivezić, *Found. Phys. Lett.* **12**, 507 (1999).
- [36] T. Aubin, *A course in Differential Geometry* (Graduate Studies in Mathematics, Vol. 27) (American Mathematical Society, Providence, 2001).